knee pain?

Your knee pain may be due to chronic swelling or inflammation in the joint – most often referred to as arthritis. Below are common forms of arthritis associated with knee pain.

With **osteoarthritis**, the cushioning cartilage at the end of the femur may have worn down, making walking painful as bone rubs against bone.²

With **rheumatoid arthritis**, sometimes called inflammatory arthritis, a person's immune system attacks the joints with uncontrolled inflammation, potentially causing joint erosion.²

With **post-traumatic arthritis**, a less common form of arthritis, a broken or fractured bone extends into the joint space, causing the surface to become uneven. Over time, friction causes the joint to break down and become arthritic.³



More than 790,000

knee replacements are performed in the United States each year⁴

Doforone

- Osteoarthritis (OA). Centers for Disease Control and Prevention. Accessed January 12, 2020. www.cdc.gov/arthritis/basics/osteoarthritis.htm
- What is arthritis? Arthritis Foundation. Accessed February 24, 2020. https://www. arthritis.org/health-wellness/about-arthritis/understanding-arthritis/what-is-arthritis
- Arthritis: an overview. OrthoInfo. Accessed April 23, 2019. https://orthoinfo.aaos.org/ en/diseases-conditions/arthritis-an-overview/
- Foran JRH. Total knee replacement. OrthoInfo. Accessed October 2020. https:// orthoinfo.aaos.org/en/treatment/total-knee-replacement/
- Hampp E, Chang TC, Pearle A. Robotic partial knee arthroplasty demonstrated greater bone preservation compared to robotic total knee arthroplasty. Poster presented at: Orthopaedic Research Society Annual Meeting; February 2-5, 2019; Austin, TX.
- Bell SW, Anthony I, Jones B, MacLean A, Rowe P, Blyth M. Improved accuracy of component positioning with robotic-assisted unicompartmental knee arthroplasty: data from a prospective, randomized controlled study. J Bone Joint Surg Am. 2016;98(8):627-635. doi:10.2106/JBJS.15.00667
- Blyth MJ, Anthony I, Rowe P, Banger MS, MacLean A, Jones B. Robotic-arm assisted versus conventional unicompartmental knee arthoplasty: exploratory secondary analysis of a randomized controlled trial. Bone Joint Res. 2017;6(11):631-639. doi:10.1302/2046-3758.611.BJR-2017-0060.R8
- Clement ND, Bell A, Simpson P, Macpherson G, Patton JT, Hamilton DF. Robotic-assisted unicompartmental knee arthroplasty has a greater early functional outcome when compared to manual total knee arthroplasty for isolated medial compartment arthritis. Bone Joint Res. 2020;9(1):15-22 doi:10.1302/2046-3758.91. BJR-2019-0147.R9
- Coon T, Shi S, DeBattista J, Bhowmik-Stoker M. Clinical and functional outcomes of robotic-arm assisted unicompartmental and bicompartmental knee arthroplasty. Presented at: European Knee Society Arthroplasty Conference; April 20-21, 2017; London, England 10.
- Vakharia RM, Law TY, Roche MW. Survivorship and patient satisfaction rates of robotic-assisted unicompartmental knee arthroplasty: a 10-year follow-up study. Presented at: American Association of Hip and Knee Surgeons (AAHKS) Annual Meeting; November 5-8, 2020; Dallas, T11.
- Alexiades MM, Boettner F, Pearle AD. Partial knee replacement: a treatment option in unicompartmental knee arthritis. Hospital for Special Surgery. Published February 13, 2013. Accessed March 29, 2018. https://www.hss.edu/conditions_partial-kneereplacement.asp

Important information

Knee replacement: Knee replacement is intended for use in individuals with joint disease resulting from degenerative, rheumatoid and post-traumatic arthritis, and for moderate deformity of the knee. Knee replacement surgery is not appropriate for patients with certain types of infections, any mental or neuromuscular disorder which would create an unacceptable risk of prosthesis instability, prosthesis fixation failure or complications in postoperative care, compromised bone stock, skeletal immaturity, severe instability of the joint, or excessive body weight.

As with any surgery, knee replacement surgery has serious risks which include, but are not limited to, pain, infection, bone fracture, peripheral neuropathies (nerve damage), circulatory compromise (including deep vein thrombosis (blood clots in the legs)), genitourinary disorders (including kidney failure), gastrointestinal disorders (including paralytic ileus (loss of intestinal digestive movement)), vascular disorders (including thrombus (blood clots), blood loss, or changes in blood pressure or heart rhythm), bronchopulmonary disorders (including emboli, stroke or pneumonia), heart attack, and death.

Implant related risks which may lead to a revision include dislocation, loosening, fracture, nerve damage, heterotopic bone formation (abnormal bone growth in tissue), wear of the implant, metal and/or foreign body sensitivity, soft tissue imbalance, osteolysis (localized progressive bone loss), and reaction to particle debris. Knee implants may not provide the same feel or performance characteristics experienced with a normal healthy joint. The information presented is for educational purposes only. Speak to your doctor to decide if joint replacement surgery is appropriate for you. Individual results vary and not all patients will return to the same activity level. The lifetime of any joint replacement is limited and varies with each individual. Your doctor will counsel you about how to best maintain your activities in order to potentially prolong the lifetime of the device. Such strategies include not engaging in high-impact activities, such as running, as well as maintaining a healthy weight. It is important to closely follow your doctor's instructions regarding post-surgery activity, treatment and follow-up care.

Ask your doctor if a knee replacement is right for you.

Stryker Corporation or its other divisions or other corporate affiliated entities own, use or have applied for the following trademarks or service marks: AccuStop, Mako, SmartRobotics, Stryker. All other trademarks are trademarks of their respective owners or holders.

MAKPKI-PE-1 Rev-1 24188

Copyright © 2021 Stryker Printed in USA



*s*tryker

325 Corporate Drive Mahwah, NJ 07430 t: 1 888 STRYKER

makosmartrobotics.com

stryker

Mako® SmartRobotics™ partial knee replacement

A patient's guide





Why do I have

To get a better idea of why your knee hurts, let's take a look at how it works. Your knee is the largest joint in your body, and it works a lot like a hinge. Three bones come together to form the joint: the lower end of the thighbone (the femur), the upper end of the shinbone (the tibia), and the kneecap (the patella) right above where the long bones meet. Tough bands called ligaments help keep everything in place and stable.

Cartilage provides cushioning, keeps bones from rubbing together, and absorbs the shock of walking, running and jumping. Your body also produces a natural lubricating fluid called synovium that minimizes friction in the joint. When everything is working smoothly, you don't have to think about the mechanics of your knee. When something's wrong, it can feel debilitating.



Over 30 million
Americans suffer from osteoarthritis¹

What is

partial knee replacement?

Partial knee replacement (PKR) is a surgical procedure that relieves pain caused by joint degeneration due to osteoarthritis (OA).

In a PKR procedure, only the damaged area of the knee joint is replaced, which may help minimize trauma to healthy bone and tissue.

There are three types of partial knee replacement:



In a **unicondylar knee replacement,** only one area (or compartment) of the joint is replaced.



In a **patellofemoral knee replacement**, the kneecap (or patella) and the groove at the lower end of the thighbone (or femur) are replaced.



In a **bicompartmental knee replacement**, two compartments of the knee are replaced—the inside (medial side) and the kneecap.

Stryker has worked with surgeons to develop innovative products like Mako SmartRobotics™ for use with partial knee replacements. With Mako SmartRobotics™, your surgeon can selectively target the part of your knee damaged by osteoarthritis.

Up to 15 percent

of all people with knee arthritis may be eligible for partial knee replacement¹¹

The Mako SmartRobotics™ difference

The difference is in the technology—AccuStop $^{\scriptscriptstyle TM}$ technology.

With Mako SmartRobotics[™], your surgeon can create a personalized surgical plan. With the help of AccuStop[™] technology, they are guided to accurately cut what's planned for you,⁶ which could help protect your healthy bone.⁵

How does Mako SmartRobotics™ and AccuStop™ technology work?







Personalized surgical plan

Before your surgery, your doctor will take a CT scan of your knee joint, which develops a 3D virtual model of your unique anatomy. This model helps your doctor see things they can't typically see with an X-ray alone—things like your bone structure and disease severity. Throughout the procedure, Mako SmartRobotics™ provides real-time data to your surgeon so they can continuously assess the movement and tension of your new joint and adjust your surgical plan if needed. Mako SmartRobotics™helps your surgeon determine the desired size, placement and positioning of your implant.

Arthritic bone removal

In the operating room, your surgeon guides Mako's robotic arm to remove arthritic bone and cartilage from the knee. AccuStop $^{\text{TM}}$ technology provides tactile resistance to help your surgeon stay within the boundaries defined in your surgical plan and accurately cut what's planned for you, 6 which could help protect your healthy bone. 5

Implant placement and range-of-motion assessment

With the removal of the diseased bone, your implant is placed into the knee joint. Once your implant is successfully placed, it's off to the recovery room to begin the journey towards strengthening your new joint.

Visit makosmartrobotics.com to download a discussion guide to use with your doctor.

Why choose Mako SmartRobotics™?

In clinical studies, compared to manual surgery, Mako SmartRobotics™ for Partial Knee resulted in:

- More accurate implant placement compared to manual surgery, which may result in improved outcomes and functioning of the knee^{6,7}
- Less pain in the days and weeks following surgery⁷
- A shorter hospital stay⁸
- Quicker recovery, where 9 out of 10 patients were walking without an aid, such as a cane or walker, three weeks after surgery⁹
- 97% of patients satisfied or very satisfied 10 years after surgery 10

